

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 3 :

C12P 21/00, 21/02  
C07H 21/04

A1

(11) International Publication Number:

WO 85/ 02200

(43) International Publication Date:

23 May 1985 (23.05.85)

(21) International Application Number: PCT/US84/01853

(22) International Filing Date: 9 November 1984 (09.11.84)

(31) Priority Application Number: 551,443

(32) Priority Date: 14 November 1983 (14.11.83)

(33) Priority Country: US

(71) Applicant: CHIRON CORPORATION [US/US]; 4560  
Horton Street, Emeryville, CA 94608 (US).(72) Inventors: BARR, Philip, J. ; 825 Lincoln Way, #301,  
San Francisco, CA 94122 (US). MERRYWEATHER,  
James, P. ; 1427 Walnut Street, Berkeley, CA 94709  
(US).(74) Agent: ROWLAND, Bertram, L.; Townsend and Town-  
send, Stewart Street Tower, One Market Plaza, San  
Francisco, CA 94105 (US).(81) Designated States: AT (European patent), BE (Euro-  
pean patent), CH (European patent), DE (European  
patent), FR (European patent), GB (European pa-  
tent), JP, LU (European patent), NL (European pa-  
tent), SE (European patent).

Published

With international search report.

(54) Title: INTERLEUKIN-2 PRODUCTION USING CLONED GENES FOR INTERLEUKIN-2 AND YEAST  
ALPHA-FACTOR

5' - IL2-6 IL2-7 IL2-8 IL2-9 IL2-10 IL2-11 IL2-12 IL2-13 - 3'

3' - IL2-19 IL2-20 IL2-21 IL2-22 IL2-23 IL2-24 IL2-25 IL2-26 IL2-27 - 5'

## (57) Abstract

Methods and compositions for efficient production of human interleukin-2. A synthetic interleukin-2 gene is joined to yeast alpha-factor secretory leader and processing signals to provide for expression and secretion of mature gene product in yeast. Enhanced yields of the product may be recovered from the nutrient medium.

8 - 448946

INTERLEUKIN-2 PRODUCTION USING CLONED GENES FOR INTERLEUKIN-2  
AND YEAST ALPHA-FACTOR

Background of the Invention

1. Field of the Invention

Lymphokines are naturally-occurring  
5 polypeptides, produced by normal lymphocytes, which  
mediate the host's immune response to antigenic  
challenge. A particular lymphokine, interleukin-2,  
appears to promote the host's immune response and has  
potential value in the treatment of tumors,  
10 immuno-deficiency diseases and several other clinical  
conditions, and as an adjuvant for vaccine  
administration. Interleukin-2 appears to act as a  
potent mitogen for T lymphocytes. At present, only  
limited quantities of interleukin-2 are obtained by  
15 separation from human serum or from certain human cell  
tissue culture media. It is therefore of great  
scientific and clinical importance to be able to  
produce sufficiently large quantities of interleukin-2.  
Economic, efficient methods for producing a product  
20 having the properties of mature human interleukin-2  
have therefore become significant goals.

2. Description of the Prior Art

Kurjan and Herskowitz, Cell (1982) 30:933-934  
describe a putative  $\alpha$ -factor precursor containing four  
25 tandem copies of mature  $\alpha$ -factor, describing the  
sequence and postulating a processing mechanism.  
Kurjan and Herskowitz, Abstracts of Papers presented at  
the 1981 Cold Springs Harbor Meeting on The Molecular  
Biology of Yeast, p. 242, in an Abstract entitled "A  
30 Putative Alpha-Factor Precursor Containing Four Tandem



Repeats of Mature Alpha-Factor," describe the sequence encoding for the  $\alpha$ -factor and spacers between two of such sequences.

- 5 Taniguchi et al., Nature (1983) 302:305-310, report the cloning of a human interleukin-2 cDNA and its expression in monkey cells in tissue culture, describe the oligonucleotide sequence and give an inferred amino acid sequence for both a putative precursor and presumed mature form. Robb et al.,  
10 (1983) Proc. Natl. Acad. Sci USA 80:5990-5994 describe a partial amino acid sequence from the N-terminal region of mature human interleukin-2 isolated from JURKAT cell tissue culture medium.

Summary of the Invention

- 15 Novel methods and DNA constructs are provided for the production of polypeptides having biological activity analogous to interleukin-2 (IL2). Enhanced efficiency in production of the polypeptide is achieved, in part, by the complete synthesis of the  
20 structural gene employing codons preferentially utilized by yeast, the intended host. Desirably, at least about 50%, usually, at least about 60% of the codons are modified such that most of the codons of the structural gene are those preferentially utilized by  
25 yeast. The construct includes a replication system for yeast and the structural gene joined in reading frame to secretory and processing signals recognized by yeast. The yeast host transformant provides for efficient and economic production of a product useful  
30 as interleukin-2.



Brief Description of the Drawings

Fig. 1 illustrates the order of assembly of synthetic ssDNA segments used in preparing the 5'-half of the synthetic IL2 gene.

5 Fig. 2 illustrates the scheme utilized to clone the synthetic DNA fragment of Fig. 1.

Fig. 3 illustrates the order of assembly of ssDNA segments used in preparing the 3'-half of the synthetic IL2 gene.

10 Fig. 4 illustrates the scheme utilized to clone the synthetic gene fragment of Fig. 3.

Fig. 5 illustrates the scheme utilized to join the two halves of the synthetic IL2 gene.

Description of the Specific Embodiments

15 DNA constructs capable of expressing mammalian, particularly human interleukin-2 (IL2) in a eukaryotic microorganism host are provided. (Unless otherwise indicated when referring to interleukin-2 (IL2) polypeptide(s), it is intended to include not  
20 only the naturally occurring mammalian factors, but also fragments or analogs thereof exhibiting analogous biological activity.) These DNA fragments can be incorporated into vectors, and the resulting plasmids used to transform susceptible hosts. Transformation of  
25 a susceptible host with such recombinant plasmids results in expression of the IL2 gene and production of a mature polypeptide product having the physiological and immunological activity of IL2. That is, it acts in the same manner as IL2 isolated from either a rat or a  
30 human host in recognized bioassays.

Extrachromosomal constructs are provided having as essential elements for expression of mature polypeptides, a replication system recognized by yeast, a synthetic structural gene having a plurality of



WO 85/02200

PCT/US84/01853

4

codons preferentially utilized by yeast, the structural gene being in reading frame with efficient secretory leader and processing signals to provide for the efficient secretion and processing of the polypeptide in a yeast host, and production of a product in high yield which has biological, particularly immunological and physiological activity commensurate with human interleukin-2. The construct provides for the initial formation of "pre"-IL2.

By "pre"-IL2, it is meant that the DNA sequence encoding the mature polypeptide is joined to and in reading frame with a leader sequence including processing signals efficiently recognized by the yeast host. Thus, "pre" denotes the inclusion of secretory leader and processing signals sequences (on the precursor polypeptide) recognized by yeast and does not refer to any processing signals associated with the natural human IL2 gene.

The IL2 structural gene of this invention is synthetic DNA, which is prepared using codons preferred by yeast as evidenced by the codon frequency in structural genes encoding for yeast glycolytic enzymes. The secretory leader and processing signals are conveniently derived from naturally-occurring DNA sequences in yeast, which sequences provide for the secretion and processing of polypeptide(s). Such polypeptides which are naturally secreted by yeast include  $\alpha$ -factor,  $\beta$ -factor, acid phosphatase, and the like. The remaining sequences in the construct, including the replication system, promoter, and terminator, are well known and amply described in the literature.

In preparing the DNA constructs of the present invention, it is necessary to bring the individual DNA sequences embodying the structural gene, the secretory leader and processing signals, replication system, promoter, and terminator together



5

in a predetermined order to assure that they are able to properly function in the resulting plasmid. Since the various DNA sequences which are joined to form the DNA construct of the present invention will be derived from diverse sources, in some instances it will be convenient or necessary to join the sequences by means of connecting or adaptor molecules, which in the subject invention are incorporated into the synthetic gene.

10 In developing the subject invention, advantage is taken of a pre-existing vector, which includes a multicopy number yeast replication system, a bacterial replication system, appropriate markers for selection, as well as a promoter, transcriptional terminator and modified leader sequence associated with the secretion of  $\alpha$ -factor. See copending application Serial No. 522,909, filed August 12, 1983.

The structural gene encoding for IL2 is prepared by first preparing a series of single stranded fragments ranging in number of from about 10 to 40, conveniently 14 to 34 bases, which provide for overlapping of other fragments and overhangs, so that upon bringing together the fragments under ligating conditions, dsDNA is produced having appropriate cohesive ends. In view of the large size of the IL2 structural gene and to provide for future flexibility in potential manipulations, a 5'- and 3'-fragment were prepared and then combined to provide for a structural gene encoding the entire amino acid sequence of IL2 and providing for appropriate termini for linking to restriction sites in the vector, which provide for the structural gene being in frame with the secretory leader sequence.

Each of the fragments encoding for the gene - 5'-fragment and 3'-fragment with flanking regions as appropriate - may be cloned and amplified in an appropriate vector to expand the amount of the fragment



and to ensure its integrity. The 5'-fragment is inserted downstream from transcriptional regulatory signals for transcription initiation and a secretory leader sequence which includes a convenient restriction site at or about the processing signal.

By employing a synthetic fragment, the termini of the fragment can be tailored to fulfill the requirements necessary for subsequent processing steps. For the 5'-fragment, its 5'-terminus is designed to join to the secretory leader sequence and processing signal in reading frame and to replace any nucleotides which have been lost due to restriction or other processing of the nucleotide sequence encoding for the secretory leader and processing signal. Thus, in restricting the vector, a restriction site can be chosen which is internal to the coding region for the secretory leader and processing signal.

In addition, it may be useful to extend the 5'-fragment in the 3'-direction beyond the site where the 5'-fragment and 3'-fragment are to be linked. In this way, a convenient 3'-terminus of the 5'-fragment is present for joining to the vector 5'-terminus. The extension is then removed by restriction to provide a 3'-terminus of the 5'-fragment which is complementary to the 5'-terminus of the 3'-fragment.

Parallel manipulations may be employed with the 3'-fragment to provide for appropriate restriction sites and termini for joining to a vector and to the 5'-fragment, as well as supplying nucleotides in the 3'-non-coding region associated with termination of transcription and translation.

After cloning the synthetic 5'-fragment in the vector containing the secretory leader sequence, a new extended 5'-fragment is excised. The new 5'-fragment has its 5'-end beginning with the transcriptional regulatory signals controlling the transcription of the secretory leader sequence and





7

processing signal, followed by the synthetic 5'-fragment in reading frame with the secretory leader sequence. Thus, this new fragment now includes the promoter and other associated transcriptional regulatory sequences, such as the TATA box and capping sequence, as well as any other sequences involved in the efficient transcription of  $\alpha$ -factor. By employment of appropriate restriction enzymes, the fragment is obtained which has all the necessary transcriptional regulatory functions, the secretory leader and processing signal, the 5'-end of the IL2 structural gene and a cohesive end or blunt end for ligating to the 5'-terminus of the 3'-fragment.

The 3'-half of the IL2 is inserted into an appropriate vector for cloning. The resulting plasmid has a unique restriction site at the 5'-end of the 3'-fragment resulting in cohesive termini when cut, and appropriate transcriptional termination sequences downstream, such as a terminator and polyadenylation signal. The fragment to be inserted conveniently has the same cohesive termini. In order to prevent circularization of the plasmid without insertion of the 5'-end of the IL2, the plasmid is treated with phosphatase. The fragment containing the transcriptional regulatory signals and 5'-end of the IL2 structural gene is then inserted into the phosphatase-treated plasmid for bacterial transformation and subsequently excised and ligated to provide for a plasmid capable of transforming a yeast host efficiently, being multicopy, and providing for the efficient secretion of the polypeptide encoded by the IL2 gene.

While the homologous promoter associated with the secretory leader sequence may be used, it may also be replaced with other promoters or may be used in tandem with other promoters.

A wide variety of promoters are available or can be obtained from yeast genes. Promoters of



particular interest include acid phosphatase and those promoters involved with enzymes in the glycolytic pathway, such as promoters for alcohol dehydrogenase, glyceraldehyde-3-phosphate dehydrogenase, pyruvate  
5 kinase, triose phosphate isomerase, phosphoglucose isomerase, phosphofructokinase, etc. By employing these promoters with regulatory sequences, such as enhancers, operators, etc., and using a host having an intact regulatory system, one can regulate the expression of  
10 the "pre"-IL2, and various small organic molecules, e.g., glucose, may be employed for the regulation of production of the desired polypeptide.

One may also employ temperature controlled systems, e.g., temperature-sensitive regulatory mutants  
15 which allow for modulation of transcription by varying the temperature. Thus, by growing the cells at either the non-permissive or permissive temperature, as appropriate, one can grow the cells to high density, before changing the temperature in order to provide for  
20 expression of the "pre"-polypeptide for IL2.

Other capabilities may also be introduced into the construct. For example, some genes provide for amplification, where upon stress to the host, not only is the gene which responds to the stress  
25 amplified, but also flanking regions. By placing such a gene upstream from the promoter, coding region and the other regulatory signals providing transcriptional control of the "pre"-polypeptide, and stressing the yeast host, plasmids may be obtained which have a  
30 plurality of repeating sequences, which sequences include the "pre"-polypeptide gene with its regulatory sequences. Illustrative genes include metallothioneins and dihydrofolate reductase.

The construct may include, in addition to the  
35 secretory leader and processing signal sequence, other DNA homologous to the host genome. If it is desired that there be integration of the IL2 gene into the



chromosome(s), integration can be enhanced by providing  
flanking sequences flanking the IL2 gene construct which are  
homologous to host chromosomal DNA.

The replication system which is employed will  
5 be recognized by the yeast host. Therefore, it is  
desirable that the replication system be native to the  
yeast host. A number of yeast vectors are reported by  
Botstein et al., Gene (1979) 8:17-24. Of particular  
interest are the YE<sub>p</sub> plasmids, which contain the 2 $\mu$ m  
10 plasmid replication system. These plasmids are stably  
maintained at multiple copy number. Alternatively, or  
in addition, one may use a combination of ARS1 and  
CEN4, to provide for stable maintenance.

The plasmids may be introduced into the yeast  
15 host by any convenient means, employing yeast host  
cells or spheroplasts and using DNA for transformation,  
or liposomes, or calcium precipitated DNA or other  
conventional technique. The modified hosts may be  
selected in accordance with the genetic markers which  
20 are usually provided in a vector used to construct the  
expression plasmid. An auxotrophic host may be  
employed, where the plasmid has a gene which  
complements the host and provides prototrophy.  
Alternatively, resistance to an appropriate biocide,  
25 e.g., antibiotic, heavy metal, toxin, or the like, may  
be included as a marker in the plasmid. Selection may  
then be achieved by employing a nutrient medium which  
stresses the parent cells, so as to select for the  
cells containing the plasmid. The plasmid containing  
30 cells may then be grown in an appropriate nutrient  
medium, and the desired secreted polypeptide isolated  
in accordance with conventional techniques. The  
polypeptide may be purified by chromatography,  
filtration, extraction, etc. Since the polypeptide  
35 will be present in mature form in the nutrient medium,  
one can cycle the nutrient medium, continuously  
removing the desired polypeptide.



10

The following examples are offered by way of example and not by way of limitation.

# EXPERIMENTAL

A nucleotide sequence for interleukin-2 comprising preferentially utilized yeast codons was devised. The sequence included a portion of a modified  $\alpha$ -factor secretory leader and processing signal at its 5'-end. The three glu-ala pairs were deleted and a base pair was changed internal to the secretory leader sequence to promote a KpnI site and change the codon from ser to gln. The sequence with the coding strand shown 5' to 3', was as follows:

## Processing Site

GlnLeuAspLysArgAlaProThrSerSerSerThrLysLysThrGlnLeuGlnLeu  
15 5'-AGCTCGGATAAAAGAGCTCCAACTCTTCCTCTACCAAGAAGCCAGCTGCAATTG  
3'-CATGTCGACCTATTTCTCGAGGTTGGAGAAGGAGATGCTTCTTGGGTCGACGTTAAC  
(KpnI)

GluHisLeuLeuLeuAspLeuGlnMetIleLeuAsnGlyIleAsnAsnTryLysAsnPro  
20 GAACACTTGTGTTGCACTTGCAAATGATCTTGAACGGTATCAACAACCTACAAGAACCCA  
CTTGTGAACAACAACCTGAACGTTTACTAGAACTTGCCATAGTTGTTGATGTTCTTGGGT

LysLeuThrArgMetLeuThrPheLysPheTyrMetProLysLysAlaThrGluLeuLys  
AAGTTGACCAGAAATGTTGACCTTCAAGTTCTACATGCCAAAGAAGGCTACCGAATTGAAG  
TTCAACTGGTCTTACAACCTGGAAGTTCAAGATGTACGGTTTCTTCCGATGGCTTAACCTC

HisLeuGlnCysLeuGluGluGluLeuLysProLeuGluGluValLeuAsnLeuAlaGln  
25 CACCTGCAGTGTCTAGACGAAGAGTTGAACCCATTGGAAGAAGTCTGAACTTGGCTCAA  
GTGGAGCTCACAGATCTCTCTCAACTTCGGTAACTTCTTCAAGGACTTGAACCGAGTT  
XbaI

SerLysAsnPheHisLeuArgProArgAspLeuIleSerAsnIleAsnValIleValLeu  
30 TCTAAGAACTTCCACTTGACACCAACAGACTTGATCTCTAATCAACGTTATCGTTTIG  
AGATTCTTCAAGGTGAACCTGTGTTCTCTGAACCTAGAGATTGTAGTTGCAATAGCAAAAC

GlnLeuLysGlySerGluThrThrPheMetCysGluTyrAlaAspGluThrAlaThrIle  
GAATTGAAGGGTTCTGAAACCACTTCATGTGTCAATACGCTGACGAAACCGCTACCATC  
CTTAACCTCCCAAGAGTTTGGTGGAGTACACACTTATCCGACTGCTTTGGCGATGGTAG

ValGluPheLeuAsnArgTrpIleThrPheCysGlnSerIleIleSerThrLeuThrOP  
35 GTTGAATTCTTCAACAGATGGATCACCTTCTGTCAATCTATCATCTCTACCTTGACCTGA  
CAACTTAAGAACTTGTCTACCTAGTGGAGACACTTAGATAGTAGAGATGGAACCTGGACT



11

AM  
TAGGCGTCG-3'  
ATCGGCAGCAGCT-5'

5                    SalI  
                  The sequence is provided with a KpnI cohesive  
end at the 5'-end and a SalI cohesive end at the  
3'-end. The coding sequence for the mature polypeptide  
begins after the processing site.

                  A synthetic DNA fragment for interleukin-2  
10 having the above sequence was prepared by synthesizing  
and cloning two halves of the fragment separately.  
Each half fragment was prepared by synthesizing  
overlapping ssDNA segments using the phosphoramidite  
method, as described by Beaucage and Caruthers (1981)  
15 Tetrahedron Lett. 22:1859-1862. The ssDNA segments  
were as follows:

| Designation | Sequence (5' to 3')                |
|-------------|------------------------------------|
| Linker      | AGCTGGATAAAAGA                     |
| IL2-1       | GCTCCAACCTCTTCCTCTACCAAGAAGACCCAG  |
| 20 IL2-2    | CTGCAATTGGAACACTTGTGTGTTGACTT      |
| IL2-3       | GCAAATGATCTTGAACGGTATCAACAACCTACA  |
| IL2-4       | AGAACCCAAAGTTGACCAGAATGTTGACCTTC   |
| IL2-5       | AAGTTCTACATGCCAAAGAAGGCTACCGAATT   |
| IL2-6       | GAAGCACCTGCAGTGTCTAGAGGAAGAGTTG    |
| 25 IL2-7    | AAGCCATTGGAAGAAGTCCTGAACTTGGCTCAAT |
| IL2-8       | CTAAGAACTTCCACTTGAGACCAAGAGACTT    |
| IL2-9       | GATCTCTAACATCAACGTTATCGTTTTGGAAT   |
| IL2-10      | TGAAGGGTTCTGAAACCACCTTGATGTGTGAA   |
| IL2-11      | TACGCTGACGAAACCGCTACCATCGTTGAAT    |
| 30 IL2-12   | TCTTGAACAGATGGATCACCTTCTGTCAATC    |
| IL2-13      | TATCATCTCTACCTTGACCTGATAGGCGTCTG   |
| IL2-14      | GAAGAGGTTGGAGCTCTTTTATCCAGCTGTAC   |
| IL2-15      | GTTCCAATTGCAGCTGGGTCTTCTTGGTAGAG   |
| IL2-16      | GTTCAAGATCATTGCAAGTCCAACAACAAGT    |
| 35 IL2-17   | GTCAACTTTGGGTCTTGTAGTTGTTGATACC    |
| IL2-18      | TTGGCATGTAGAACTTGAAGGTCAACATTCTG   |
| IL2-19      | ACACTGCAGGTGCTTCAATTCGGTAGCCTTCT   |
| IL2-20      | TTCTTCCAATGGCTTCAACTCTTCCTCTAG     |



WO 85/02200

PCT/US84/01853

12

|          |                                     |
|----------|-------------------------------------|
| IL2-20'  | TCGACAACTCTTCCTCTAG                 |
| IL2-21   | AGTGGAAAGTTCTTAGATTGAGCCAAGTTCAGGAC |
| IL2-22   | GTTGATGTTAGAGATCAAGTCTCTTGGTCTCA    |
| IL2-23   | GTTTCAGAACCCTTCAATTCCAAAACGATAAC    |
| 5 IL2-24 | GGTTTCGTCAGCGTATTACACATGAAGGTG      |
| IL2-25   | TCCATCTGTTCAAGAATTCAACGATGGTAGC     |
| IL2-26   | AAGGTAGAGATGATAGATTGACAGAAGGTGA     |
| IL2-27   | TCGACGACGCCTATCAGGTC                |

The 5'-half of the sequence was assembled as illustrated in Fig. 1. Fifty pmoles of each ssDNA segment (except Linker and IL2-20') were 5'-phosphorylated with T4 polynucleotide kinase. The segments were then annealed in a single step by combining and cooling from 95°C to 25°C over 1.5 hours. Ligation was performed in a reaction volume of 30 µl containing 1mM ATP, 10mM DTT, 100mM tris-HCl, pH 7.8, 10mM MgCl<sub>2</sub>, 1µg/ml spermidine and T4 ligase. The resulting double stranded fragment was purified on a 7% native polyacrylamide electrophoresis gel. The dsDNA fragment included a KpnI cohesive end at the 5'-end and a SalI cohesive end at the 3'-end.

After assembly, the 5'-half of the sequence was inserted into pEGF-24 downstream and in frame with the modified  $\alpha$ -factor secretory leader and processing signal. The plasmid pEGF-24 is described in application serial no. 522,909, filed August 12, 1983, which pertinent portion is incorporated by reference and reproduced in pertinent part as follows.

A synthetic sequence for human epidermal growth factor (EGF) based on the amino acid sequence of EGF reported by H. Gregory and E. M. Preston Int. J. Peptide Protein Res. 7, 107-116 (1977) was prepared. The sequence was inserted into the EcoRI site of pBR328 to produce a plasmid p328EGF-1 and cloned.

Approximately 30µg of p328EGF-1 was digested with EcoRI and approximately 1µg of the expected 190



Herskowitz, Abstracts 1981 Cold Spring Harbor meeting on the Molecular Biology of Yeasts, page 242).

The resulting mixture was used to transform E. coli HB101 cells and plasmid pAB201 obtained.

- 5 Plasmid pAB201 (5ug) was digested to completion with the enzyme EcoRI and the resulting fragments were:
- a) filled in with DNA polymerase I Klenow fragment;
  - b) ligated to an excess of BamHI linkers; and
  - c) digested with BamHI. The 1.75 kbp EcoRI fragment
- 10 was isolated by preparative gel electrophoresis and approximately 100ng of the fragment was ligated to 100ng of pC1/1, which had been previously digested to completion with the restriction enzyme BamHI and treated with alkaline phosphatase.
- 15 Plasmid pC1/1 is a derivative of pJDB219, Beggs, Nature (1978) 275:104, in which the region corresponding to bacterial plasmid pMB9 in pJDB219 has been replaced by pBR322 in pC1/1. This mixture was used to transform E. coli HB101 cells. Transformants
- 20 were selected by ampicillin resistance and their plasmids analyzed by restriction endonucleases. DNA from one selected clone (pYEGF-8) was prepared and used to transform yeast AB103 cells. Transformants were selected by their leu<sup>+</sup> phenotype.
- 25 The above described construction was modified using different sequences for joining structural genes to the  $\alpha$ -factor secretory leader sequence and/or site specific mutagenesis, thus providing for different processing signals. In the following table, a. through
- 30 e. show the sequence of the fusions at the N-terminal region of the structural gene hEGF, as exemplary, which sequences differ among the several constructions.



15

80 Gly Val Ser Leu Asp Lys Arg Glu Ala Glu Ala Glu Ala Asn Ser Asp Ser Glu 5  
85 /  
GGG GTA TCT TTG CAT AAA AGA GAG GGT GAA GCT AAC TCC GAC TCC GAA  
CCC CAT AGA AAC CTA TTT TCT CTC CGA CTT CGA TTT CGA TTG AGG CTG AGG CTT  
80 85 90  
Gly Val Ser Leu Asp Lys Arg Glu Ala Glu Ala Ser Leu Asp Lys Arg Asn Ser Asp Ser Glu 5  
85 /  
GGG GTA TCT TTG CAT AAA AGA GAG GGT GAA GCT AAC TCC GAC TCC GAA  
CCC CAT AGA AAC CTA TTT TCT CTC CGA CTT CGA TTT TCT TCT TTG AGG CTG AGG CTT  
80 85 90  
Gly Val Ser Leu Asp Lys Arg Asn Ser Asp Ser Glu 5  
85 /  
GGG GTA TCT TTG CAT AAA AGA GAG GGT GAA GCT AAC TCC GAC TCC GAA  
CCC CAT AGA AAC CTA TTT TCT CTC CGA CTT CGA TTT TCT TCT TTG AGG CTG AGG CTT  
80 85 90  
Gly Val Pro Leu Asp Lys Arg Asn Ser Asp Ser Glu 5  
85 /  
GGG GTA TCT TTG CAT AAA AGA GAG GGT GAA GCT AAC TCC GAC TCC GAA  
CCC CAT AGA AAC CTA TTT TCT CTC CGA CTT CGA TTT TCT TCT TTG AGG CTG AGG CTT  
80 85 90  
Gly Val Glu Leu Asp Lys Arg Asn Ser Asp Ser Glu 5  
85 /  
GGG GTA TCT TTG CAT AAA AGA GAG GGT GAA GCT AAC TCC GAC TCC GAA  
CCC CAT AGA AAC CTA TTT TCT CTC CGA CTT CGA TTT TCT TCT TTG AGG CTG AGG CTT  
80 85 90  
Tyr Trp Glu Leu Arg 53  
Linker-4  
TCC TCC GAA TGG AGA TGA TAA GTC GAC CGA TG  
ACC ACC CTT AAC TCT ACT ATT CAG CTG GCT AC

(pYpEGF-21)

(pYpEGF-22)

(pYpEGF-23)

(pYpEGF-24)

(pYpEGF-25)

SUBSTITUTE SHEET





16

F. shows the sequences at the C-terminal region of hEGF, which is the same for all constructions. Synthetic oligonucleotide linkers used in these constructions are boxed.

5        These fusions were made as follows.

Construction (a) was made as described above. Construction (b) was made in a similar way except that linker 2 was used instead of linker 1. Linker 2 modifies the  $\alpha$ -factor processing signal by inserting an additional processing site (ser-leu-asn-lys-arg) immediately preceding the hEGF gene. The resulting yeast plasmid is named pY $\alpha$ EGF-22. Construction (c), in which the dipeptidyl aminopeptidase maturation site (glu-ala) has been removed, was obtained by in vitro mutagenesis of construction (a). A PstI-SalI fragment containing the  $\alpha$ -factor leader-hEGF fusion was cloned in phage M13 and isolated in a single-stranded form. A synthetic 31-mer of sequence

5'-TCTTTGGATAAAAGAAACTCCGACTCCCG-3'

20        was synthesized and 70 picomoles were used as a primer for the synthesis of the second strand from 1 picomole of the above template by the Klenow fragment of DNA polymerase. After fill-in and ligation at 14° for 18 hrs., the mixture was treated with  $S_1$  nuclease (5 units for 15 minutes) and used to transfect E. coli JM101 cells. Bacteriophage containing DNA sequences in which the region coding for (glu-ala) was removed were located by filter plaque hybridization using the  $^{32}$ P-labelled primer as probe. RF DNA from positive 30        plaques was isolated, digested with PstI and SalI and the resulting fragment inserted in pAB114 which had been previously digested to completion with SalI and partially with PstI and treated with alkaline phosphatase.

35        The plasmid pAB114 was derived as follows: plasmid pAB112 was digested to completion with HindIII and then religated at low (4 $\mu$ g/ml) DNA concentration



17

and plasmid pAB113 was obtained in which three 63bp  
HindIII fragments have been deleted from the  $\alpha$ -factor  
structural gene, leaving only a single copy of mature  
 $\alpha$ -factor coding region. A BamHI site was added to  
5 plasmid pAB11 by cleavage with EcoRI, filling in of the  
overhanging ends by the Klenow fragment of DNA  
polymerase, ligation of BamHI linkers, cleavage with  
BamHI and religation to obtain pAB12. Plasmid pAB13  
was digested with EcoRI, the overhanging ends filled  
10 in, and ligated to BamHI linkers. After digestion with  
BamHI the 1500bp fragment was gel-purified and ligated  
to pAB12 which had been digested with BamHI and treated  
with alkaline phosphatase. Plasmid pAB114, which  
contains a 1500bp BamHI fragment carrying the  $\alpha$ -factor  
15 gene, was obtained. The resulting plasmid (pAB114  
containing the above described construct) is then  
digested with BamHI and ligated into plasmid pCI/1.

The resulting yeast plasmid is named  
pY $\alpha$ EGF-23. Construction (d), in which a new KpnI site  
20 was generated, was made as described for construction  
(c) except that the 36-mer oligonucleotide primer of  
sequence 5'-GGGTACCTTTGGATAAAAGAACTCCGACTCCGAAT-3' was  
used. The resulting yeast plasmid is named pY $\alpha$ EGF-24.  
Construction (e) was derived by digestion of the  
25 plasmid containing construction (d) with KpnI and SalI  
instead of linker 1 and 2. The resulting yeast plasmid  
is named pY $\alpha$ EGF25.

The 5'-half of the IL2 sequence was inserted  
according to the scheme illustrated in Fig. 2. Plasmid  
30 p $\alpha$ EGF-24 was restricted with a mixture of restriction  
endonucleases KpnI and SalI to remove a KpnI/SalI  
fragment. The 5'-synthetic fragment of IL2 was  
inserted into the resulting cut vector to produce  
plasmid p $\alpha$ IL2-5' which was then cloned in E. coli  
35 HB101.

The 3'-half of the sequence was assembled as  
illustrated in Fig. 3. Fifty pmoles of each ssDNA



18

segment (except IL2-6 and IL2-27) were  
5'-phosphorylated with T4 polynucleotide kinase. The  
segments were then annealed in a single step by  
combining and cooling from 95°C to 25°C over 1.5 hours.  
5 Ligation was performed in a reaction volume of 30 µl  
containing 1mM ATP, 10mM DTT, 100mM tris-HCl, pH 7.8,  
10mM MgCl<sub>2</sub>, 1 µg/ml spermidine and T4 ligase. The  
resulting double stranded fragment was purified on a 7%  
native polyacrylamide electrophoresis gel. The dsDNA  
10 fragment included a XbaI cohesive end at the 5'-end and  
a SalI cohesive end at the 3'-end.

After assembly, the 3'-half of the sequence  
was inserted in XbaI/SalI digested pAB114. Plasmid  
pAB114 is described in application serial no. 522,909  
15 which has in part been reproduced above.

The 3'-half of the sequence was cloned  
according to the scheme illustrated in Fig. 4. Plasmid  
pAB114 was restricted with a mixture of restriction  
endonucleases XbaI and SalI to remove a XbaI/SalI  
20 fragment. The 3'-synthetic fragment was inserted into  
the resulting cut vector to produce pαIL2-3' which was  
then cloned in E. coli HB101.

After amplification, the two halves of the  
synthetic IL2 sequence in frame with yeast secretory  
25 and processing signals, as well as joined at the 5'-end  
to yeast transcriptional regulatory signals, were  
joined together in the pαIL2-3' plasmid according to  
the scheme illustrated in Fig. 5. Plasmid pαIL2-5' was  
restricted with restriction endonuclease XbaI to  
30 generate an XbaI/XbaI fragment carrying the α-factor  
transcriptional regulatory sequences and modified  
secretory leader and processing signals derived from  
plasmid pαEGF-24 and the 5'-half of the IL2 sequence.  
The 3'-proximal XbaI site in the 5'-segment is located  
35 interior of the synthetic sequence so that a 15 bp  
segment at the 3'-end is removed. Plasmid pαIL2-3' was



WO 85/02200

PCT/US84/01853

20

bioassay for human interleukin-2. A semi-quantative assay was employed by visually estimating cell survival and growth employing a microscop and comparing test results with standards employing known amounts of IL2.

5 Serial 2-fold dilutions of the yeast medium dialysate above were then prepared in mammalian cell growth medium (RPMI-1640 with supplements, as above) and 100  $\mu$ l of each diluted sample added to individual wells containing HT-2 cells.

10 Reference standards known to contain interleukin-2 (concanavalin A-free, conditioned rat splenocyte medium) were either obtained commercially (Monoclone<sup>®</sup>, Collaborative Research, Inc.) or prepared by stimulation of rat spleen cell cultures (in  
15 RPMI-1640 medium with supplements, as above) with concanavalin A (1  $\mu$ g/ $10^6$  cells) for 48 hours at 37°C in 7% CO<sub>2</sub>/air, followed by recovery of the medium, absorption with Sephadex<sup>®</sup> G-25 to remove concanavalin A and filter sterilization. A 2-fold dilution series of  
20 each reference standard was prepared as described for the yeast medium dialysate.

The HT-2 cell microplate cultures were incubated at 37°C in 7% CO<sub>2</sub>/air for 48 hours, scored for viability and/or growth and the approximate  
25 interleukin-2 content of the yeast preparation determined by reference to the standards. This comparison indicated activities equivalent to or greater than the commercial material, i.e., estimated to be in the range of 20-100 ng/ml, probably about 50  
30 ng/ml.

In accordance with the subject invention, novel DNA constructs are provided which may be inserted into vectors to provide for expression of "pre"-interleukin-2 and intracellular processing and  
35 secretion of the mature polypeptide in good yield to promote a polypeptide product having high IL2 biological activity in a recognized bioassay based on



murine cellular growth. Thus, it is possible to obtain a polypeptide having the physiological activity of the naturally-occurring human interleukin-2. By providing for secretion, greatly enhanced yields can be obtained and subsequent isolation and purification of the polypeptides simplified.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.



WO 85/02200

PCT/US84/01853

22

## WHAT IS CLAIMED IS:

1. A method for producing a biologically active polypeptide having the biological activity of human IL2 in good yield, said method comprising:
- growing yeast transformants having a
- 5 multicopy extrachromosomal element including a DNA construct which comprises in the direction of transcription, yeast recognized transcriptional regulatory sequences, and secretory leader and
- 10 and processing signals a synthetic gene encoding at least substantially for the polypeptide sequence of human IL2, stop codon(s) and a terminator, whereby a mature polypeptide having at least substantially the same amino acid sequence of human IL2 is secreted; and
- 15 isolating said mature polypeptide from the medium having the biological activity of human IL2.

2. A method according to Claim 1, wherein the coding strand of said synthesized sequence has the following nucleotide sequence:

20 GCTCCAACTCTTCTCTACCAAGAAGACCCAGCTGCAATTG  
GACTTGTGTGCACTTGCAAATGATCTTGACGGTATCAACAACCTACAAGAACC  
AAGTTGACCAGAAATGTTGACCTTCAAGTTCTACATGCCAAAGAAGCTACCGAATTGAAG  
CACCTGCAGTGTCTACAGGAACAGTTGAAGCCATTGGAAGAAGTCTGAACTTGGCTCAA  
TCTAAGAACTTCCACTTGAGACCAAGAGACTTGATCTCTAACATCAACGTTATGTTTTG  
25 GAAATGAAGGGTCTGAAACCAGCTTCATGTGIGAATACGCTGACGAAACCGCTACCATC  
GTTGAAITCTTGAACAGATGGATCACCTTCTGTCAATCTATCATCTCTACCTTGACC

3. A method according to Claim 2, wherein said secretory leader and processing signal is derived from g-factor.

- 30 4. A method according to Claim 3, wherein said g-factor secretory leader and processing signal



23

ar modified at least by removal of the glu-ala dipeptide codons.

5. A method according to Claim 3, wherein said DNA sequence is synthesized by producing two or more dsDNA fragments with flanking regions for linking to other sequences, which fragments are joined together to produce the entire sequence.

6. A DNA construct comprising in the direction of transcription yeast recognized transcriptional regulatory signals and a secretory leader sequence and processing signal; in reading frame with said secretory leader and processing signals, a synthetic DNA sequence encoding for the human IL2 gene having at least a plurality of codons preferentially utilized by yeast; stop codon(s); and, a transcriptional terminator.

7. A DNA construct according to Claim 6, wherein said DNA sequence is as follows:

GCTCCAACCTCTTCTCTACCAAGAAGACCCAGCTGCAATTG  
20 GAACACTTGTTGTGGACTTGTAAATGATCTTGAACCGTATCAACAATAAAGAACCCA  
AAGTTCACCAGAAATGTTGACCTTCAAGTTCTACATGCCAAAGAAGGCTACCGAATTGAAG  
CACCTGCAGTGTCTAGAGGAAGAGTTGAAGCCATTGCAAGAACTCCTGAACTTGGCTCAA  
TCTAAGAACTTCCACTTCAGACCAAGAGACTTGATCTCTAACATCAACGTTATCGTTTTG  
GAATTGAAGGGTTCTGAAACCACCTTCATGTGTCAATACGCTGACGAAACCGCTAGCATC  
25 GTTGAATTCTTGAACAGATGCATCACCTTCTGTCAATCTATCATCTCTACCTTGACC

8. A DNA construct according to Claim 7, wherein said secretory leader and processing signal is derived from the g-factor secretory leader and processing signal.



24

9. A DNA construct according to Claim 8, wherein said g-factor secretory leader and processing signal are modified at least by removal of the glu-ala dipeptide codons.





1/3




FIG. 1.




FIG. 2.


SUBSTITUTE SHEET



2 / 3

5'- IL2-6 IL2-7 IL2-8 IL2-9 IL2-10 IL2-11 IL2-12 IL2-13 -3'  
3'- IL2-19 IL2-20 IL2-21 IL2-22 IL2-23 IL2-24 IL2-25 IL2-26 IL2-27 -5'

FIG. 3.



XbaI SalI

FIG. 4.

SUBSTITUTE SHEET



WO 85/02200

3/3




FIG. 5.

SUBSTITUTE SHEET

